

The impact of abuse and neglect on the health and mental health of children and young people

Anne Lazenbatt
NSPCC Reader in Childhood Studies, Queen's University Belfast

February 2010

Summary

During the past 30 years, the focus on the extent and nature of child abuse and neglect has been coupled with an increasing interest in the impact on children's development, health and mental wellbeing. Child maltreatment is both a human rights violation and a complex public health issue, likely caused by a myriad of factors that involve the individual, the family, and the community. Child abuse includes any type of maltreatment or harm inflicted upon children and young people in interactions between adults (or older adolescents). Such maltreatment is likely to cause enduring harm to the child.

The different forms of abuse and neglect often occur together in one family and can affect one or more children. These include, in decreasing level of frequency: neglect; physical abuse and non-accidental injury; emotional abuse; and sexual abuse (Cawson et al, 2000; 2002). Recently, bullying and domestic violence have been included as forms of abuse of children.

There is a sizeable body of literature on the relationship between types of child maltreatment and a variety of negative health and mental health consequences. These include biological, psychological, and social deficits (for reviews, see Crittenden, 1998; Kendall-Tackett, 2001; 2003). Aside from the serious physical and health consequences of child maltreatment, several emotional and behavioural consequences for children have been noted in the literature.

These consequences vary according to differences in the severity, duration, and frequency of maltreatment. However, they also vary depending on the child's resilience, which relates to temperament, coping skills, and developmental stage, and his or her environment, as determined by family income, social support, or neighbourhood characteristics (Hecht and Hansen, 2001). Sustained maltreatment can have major long-term effects on all aspects of children's health and wellbeing.

Key findings

- Evidence states that the experience of maltreatment can have major long-term effects on all aspects of a child's health, growth and intellectual development and mental wellbeing and that it can impair their functioning as adults.
- The impact of child maltreatment includes a wide range of many complex social and economic problems, with an increased likelihood of mental disorders, health problems, education failure and unemployment, substance addiction, crime and delinquency, homelessness and an intergenerational cycle of abuse and neglect.
- The health effects of child abuse include physical injuries such as shaken baby syndrome, non-organic failure to thrive, broken bones, spinal injuries, stomach aches, migraines, and gut problems. Health problems later in life can include heart disease, obesity, liver disease, cancer and chronic lung disease.
- Depression, severe anxiety, panic attacks and post-traumatic stress disorder (PTSD) are the most common mental health consequences of abuse: the literature suggests that between 30 and 50 per cent of sexually abused children meet the full criteria for a PTSD diagnosis (Widom 1999; Darves-Bornoz et al. 1998), and up to 80 per cent experience at least some 'post-traumatic' symptoms (McLeer et al, 1992; Cuffe et al, 1998). These symptoms include hyper-vigilance, intrusive thoughts, and sudden intrusive flashbacks of the abuse experience.

Research briefing

- The impact of child maltreatment is often described as physical, psychological, behavioural, or societal. In reality, however, it is unrealistic to view these consequences in isolation. Depression and anxiety, for example, may make a young person more likely to smoke, abuse alcohol or illicit drugs, or overeat. High-risk behaviours, in turn, can lead to long-term physical health problems such as sexually transmitted diseases, drug and alcohol addiction, cancer, and obesity.
- Child physical abuse is associated with a wide range of debilitating emotional and behavioural problems that may persist into adulthood and generalize to future relationships, including parent-child relationships. It can lead directly to neurological damage, physical injuries, pain and disability or, in extreme cases, death. It has been linked to aggressive behaviour, emotional and behavioural problems, and educational difficulties in children (Finkelhor, 2008).
- Emotional abuse has an important impact on a developing child's mental health, behaviour and self-esteem. It can be especially damaging during the critical period of infancy, and affect children especially during their school years.
- Persistent neglect can lead to serious impairment of health and development, and long-term difficulties with social functioning, relationships and educational progress. In extreme cases, neglect can also result in death (Sidebotham, 2007).
- Sexual abuse is linked to disturbed mental health resulting in self-harm, inappropriate sexualised behaviour, sadness, depression and loss of self-esteem. These adverse effects may endure into adulthood.
- Research by Sidebotham (2007) suggests that up to 40 per cent of maltreatment-related deaths are probably the result of neglect, or a combination of neglect and other forms of abuse, with death resulting from extreme malnutrition, electrolyte imbalance, hypothermia, or infection. Many fatal cases seem to have an element of intent to deprive the child of his or her needs.

Research briefing

- Domestic abuse can directly and indirectly affect children, even before birth. It is likely to have a damaging effect on their health and development, with children under one at the highest risk of injury or death (McVeigh et al. 2005; Goodall and Lumley 2007).
- Domestic abuse, adult mental ill-health problems, substance misuse or racism from a caregiver may be factors underlying the physical, sexual, and emotional abuse of children within the family.
- For children with disabilities, the usual risk factors for child abuse (i.e. dependence and vulnerability), are intensified. When a child or young person is disabled, injuries or behavioural symptoms can mistakenly be attributed to the disability rather than the abuse or neglect.
- Abused children follow several distinct developmental trajectories (Noll et al, 2003; 2006). For example, victims suffering seemingly more mild forms of abuse can appear to be asymptomatic in the acute phases immediately following disclosure, but may become more symptomatic later in development (Trickett and Putman, 1998). These findings suggest that treatment of child abuse should either continue throughout development, or be revisited when issues reminiscent of the abuse become developmentally salient.

Introduction: effects of maltreatment on children's health

In many cases, child maltreatment has consequences for children, families, and society that last lifetimes (Kendall-Tackett, 2003). Infants and young children are particularly vulnerable to the physical effects of maltreatment.

Physical abuse is associated with various types of injuries, particularly when exposure to such abuse occurs in the first three years of life (Vinchon et al, 2005). Shaking an infant may result in bruising, bleeding, and swelling in the brain. The physical consequences of 'shaken baby syndrome' can range from vomiting or irritability to more severe effects, such as concussions, respiratory distress, seizures, and death (Conway, 1998). Two-thirds of subdural haemorrhages in children under two are caused by physical abuse (Vinchon et al, 2005). It is estimated that 10 per cent of admissions to paediatric burns and plastic surgery units are related to child maltreatment (Chester et al, 2006).

Infants who have been neglected and malnourished may also experience a condition known as 'non-organic failure to thrive'. This refers to a situation in which the child's weight, height, and motor development fall significantly below age-appropriate ranges, without a medical or organic cause. In extreme cases, the death of the child is the end result. Even with treatment, the long-term consequences can include continued growth problems, retardation, and socio-emotional deficits (Wallace, 1996).

Domestic abuse poses a serious risk even to the unborn foetus, as violence may increase the risk of premature birth, low birth weight, chorioamnionitis, foetal injury and in the worse case, death (Mezey and Bewley 1997, Connolly et al 1997, Bacchus et al 2002). It has been suggested that foetal morbidity resulting from violence is more prevalent than that from gestational diabetes or pre-eclampsia (Sidebotham and Golding, 2001). Foetal abuse can have effects on the developing infant's brain, leading to childhood anxiety and hyperactivity (Hosking and Walsh, 2005).

Research briefing

New technologies such as functional MRI (magnetic resonance imaging) and PET (positron emission tomography) have enabled scientists to identify the chemical and structural differences between the central nervous systems of abused and non-abused young people (Anderson et al, 2002; Teicher et al, 2004; Weniger et al, 2008). Many health problems, including panic or post-traumatic stress disorder, chronic fatigue syndrome, fibromyalgia, depression, some auto-immune disorders, suicidal tendencies, abnormal fear responses, pre-term labour, chronic pain syndromes, and ovarian dysfunction can be understood, in some cases, as manifestations of childhood maltreatment (Kendall-Tackett, 2000; De Bellis, 2005).

Evidence shows that maltreatment may inhibit the appropriate development of certain regions of the brain (Glaser, 2000). A neglected infant or young child may not be exposed to stimuli that normally activate important regions of the brain and strengthen cognitive pathways. The connections among neurons in these inactivated regions can literally wither away, hampering the child's functioning later in life. As a result, the brain may become 'wired' to experience the world as hostile and uncaring. This negative perspective may influence the child's later interactions, prompting the child to become anxious and overly aggressive or emotionally withdrawn.

Neglect and other forms of abuse may also be associated with neuromotor handicaps, such as central nervous system damage, physical defects, growth and mental retardation, and speech problems (Chester, 2006). Recent studies have also found an association between childhood abuse and hormonal disruption, manifesting in a dysregulation of the HPA (hypothalamic pituitary adrenal) axis (Cicchetti and Rogosch, 2001). In addition, childhood abuse also has strong links to later health problems, including heart disease, liver disease, cancer and chronic lung disease (Felitti et al, 1998).

Maltreatment may affect a child's health indirectly. For instance, physical and sexual abuse is a major factor in the homelessness of young people, which may result in risk-taking behaviours including substance abuse, self-harming, prostitution, and increased vulnerability

to further assault. Child victims of sexual abuse, for example, may be more prone to sexually transmitted infections, including syphilis and HIV (human immunodeficiency virus).

Importantly, abnormal ano–genital signs are uncommon in children examined for suspected child sexual abuse (Heger et al, 2002; RCPCH, 2007). Adolescents who have experienced sexual abuse are more likely to experience ongoing health problems such as chronic pelvic pain and other gynaecologic problems, gastrointestinal problems, headaches, and increased obesity (Springer et al, 2007). Both physical and sexual abuse are associated with a doubling of the risk of attempted suicide for young people by the time they reach their late twenties (Gilbert et al, 2008).

The link between maltreatment and many of these adverse consequences may be stress and depression, which can influence the immune system and may lead to higher risk-taking behaviours such as smoking, abuse of alcohol, illegal drugs, and overeating (Widom and Maxfield, 2001).

The broad range of direct and indirect health effects of child maltreatment is likely to have a substantial impact on a victim's life expectancy and long-term health-related quality of life (HRQL).

Effects on children's mental health and wellbeing

All types of maltreatment can affect a child's emotional, psychological and mental wellbeing, and these consequences may appear immediately or years later. The immediate and longer-term impact of abuse can include mental health problems such as anxiety, depression, substance misuse, eating disorders, self-injurious behaviour, anger and aggression, sexual symptoms and age-inappropriate sexual behaviour (Lanktree et al, 2008).

Numerous studies have documented associations between a child's exposure to maltreatment with negative mental health outcomes: low self-esteem and depression (Briere, 1996; Heim

and Nemeroff, 2001); severe anxiety (Kendler et al, 1998); addictions, drug and alcohol abuse (Bremner et al, 2000); post-traumatic stress disorder (McCauley et al, 1997); self-harming and suicidality (Oates, 2003); and being bullied (Duncan, 1999).

Other psychological and emotional conditions include panic disorder, dissociative disorders, attention deficit/hyperactivity disorder, and reactive attachment disorder (Teicher, 2000; De Bellis and Thomas, 2003; Springer et al, 2007). In one long-term study by Silverman et al (1996), as many as 80 per cent of young adults who had been abused met the diagnostic criteria for at least one psychiatric disorder by the time they reached age 21. These young adults exhibited many problems, including depression, anxiety, eating disorders, and suicide attempts.

Children who experience rejection or neglect are more likely to develop antisocial traits as they grow up and are more associated with borderline personality disorders and violent behaviour (Schore, 2003). Abused and neglected adolescents are estimated to be at least 25 per cent more likely to experience problems such as delinquency, teen pregnancy, low academic achievement, drug use, and mental health problems (Kelley et al 1997). Other studies suggest that abused young people are likely to engage in sexual risk-taking as they reach adolescence, thereby increasing their chances of contracting sexually transmitted infections (Johnson et al, 2006).

Evidence shows that around 50 per cent of people receiving mental health services report abuse as children: one review found that “on careful questioning, 50 to 60 per cent of psychiatric inpatients and 40 to 60 per cent of outpatients report childhood histories of physical or sexual abuse or both.” (Read, 1998). Others have concluded that “child abuse may have a causative role in the most severe psychiatric conditions.” (Fergusson et al, 1996; Mullen et al, 1993).

While the negative effects on health and development can often, though not always be reversed, this requires timely identification of the maltreatment and appropriate intervention.

The harmful effects vary depending on a number of factors, including the circumstances, personal characteristics of the child, and the child's environment (Gelles, 1998), and may endure long after the abuse or neglect occurs. Researchers have identified links between child maltreatment with difficulties during infancy, such as depression and withdrawal symptoms, common among children as young as three who have experienced emotional or physical abuse, or neglect (Dubowitz et al, 2002). Heim and Nemeroff (2001) suggest that early childhood abuse and trauma can cause a persistent biological state, which is likely to function as a risk factor for the occurrence of mental disorders in later life. It follows that abuse in childhood should be recognised as an important risk factor for mental disorders (Agid et al, 2000). Persistent neglect can lead to serious impairment of health and development; children may also experience low self-esteem or feelings of being unloved and isolated.

Domestic Abuse and its Effects on Children

Research has consistently shown that a high proportion of children living with domestic violence are themselves being abused, either physically or sexually, by the same perpetrator. Walby and Allen (2004) report a co-occurrence of domestic violence and child abuse in 40 per cent of cases, while Mullender et al (2003, 2005) estimate that in 90 per cent of incidents, children are witnesses to the violence.

Prolonged and/or regular exposure to domestic abuse can, despite the best efforts of the parents to protect the child, seriously affect the child's development, health and emotional wellbeing in a number of ways. It poses a threat to unborn children (Bacchus et al 2002), because assaults on pregnant women frequently involve punches or kicks directed at the abdomen, risking injury to both mother and foetus (Jasinski, 2004).

Domestic abuse during pregnancy and the first six months of child rearing is significantly related to various types of child maltreatment (child physical abuse, neglect, and emotional abuse) up to the child's fifth year, with children under one year at the highest risk of injury or death (Goodall and Lumley 2007). Older children may also suffer blows during episodes of

violence, and children who live in homes where domestic violence occurs are 15 times more likely to be physically abused or seriously neglected compared to the general child population (Carlson, 2000).

Children may also be greatly distressed by witnessing the physical and emotional suffering of a parent (Mullender, 2005; Hester et al. 1998; McGee 2000; Mullender et al. 2003), which can in itself be psychologically and emotionally harming. Studies by Silvern et al (1995) and Singer et al (1998) indicate that child witnesses to domestic violence are, on average, more aggressive and fearful and more often suffer from severe anxiety, depression and other trauma-related symptoms. They live with constant anxiety and may be at a higher risk of alcohol or drug abuse, experience cognitive problems or stress-related ailments (headaches, rashes), and have difficulties in school.

Multiple adverse events in childhood

Rosen and Martin (1996) have drawn attention to the fact that research into child abuse often focuses on only one type of abuse. This, however, overlooks the combined effect of different types of abuse: Horwitz's prospective study (2003), for instance, suggests that children who have experienced child sexual abuse often grow up in impoverished environments, with poverty, inadequate parenting, parents who are unemployed, or parents using drugs or alcohol. Such children have often experienced other forms of child maltreatment as well, including emotional abuse, neglect, physical abuse, and witnessing domestic abuse in the home (Coid et al, 2001; Radford and Hester, 2006). About one-third of adults self-report that they have experienced more than one form of child maltreatment (Edwards et al, 2003). Indeed, some researchers have suggested that emotional abuse is inherent in all forms of maltreatment and cannot be disentangled from other types of abuse (Garbarino et al, 1986). Emotional abuse can have a severe impact on a developing child's mental health, behaviour and self-esteem, particularly when it occurs in infancy. Underlying emotional abuse may be as important, if not more so, than other, more visible forms of abuse, in terms of its impact on the child (Glaser et al, 2001).

The effect of the co-occurrence of multiple categories of maltreatment on psychological health and wellbeing has often been overlooked, although one study by Felitti et al (2002) indicates that the effects are negatively related to the number of abuse types experienced. The study, which examined adult patients of an American health maintenance organisation, also reported a negative/positive dose-response relationship between the number of indicators of childhood maltreatment or family dysfunction, and a broad range of health outcomes. That is, as the number of negative experiences increased, poorer health was reported.

Protective Factors and Resilience

The past 20 years of research have brought an awareness of the vast individual differences in acute and long-term responses to childhood abuse. Studies have shown a relationship between various forms of childhood abuse and poor health (Flaherty et al, 2008; Felitti, 2002). However, in some cases, children may not appear to exhibit significant effects from maltreatment. This may be because they have certain protective qualities and are more resilient to negative consequences, buffered by personal characteristics such as optimism, high self-esteem or a sense of hopefulness despite their circumstances. Furthermore, there are individual differences in the timing of manifesting symptoms; some victims display few symptoms initially but evidence ‘sleeper’ effects¹ later in development (Finkelhor and Baron, 1986; Trickett and Putman, 1998).

Methodological Issues

The process by which maltreatment leads to negative health outcomes, including the causal role of maltreatment, is not fully understood. This is primarily because of the lack of well-developed theory and methodologically rigorous studies that examine factors such as poor

¹ The ‘sleeper effect’ is a psychological phenomenon whereby a highly persuasive message or event (such as child abuse), paired with a discounting cue, causes an individual to be more rather than less persuaded by the message over time: time does not heal but rather adds to the symptomatology.

Research briefing

attachment, poor parenting, poverty, drug and alcohol abuse, mental illness, and how they relate to child maltreatment. Many studies have employed cross-sectional designs, comparing the health states of individuals who report child maltreatment with those who do not (Felitti et al, 1998). More longitudinal studies are needed to better examine the processes by which maltreatment leads to negative outcomes.

Practice implications

1. Individual practitioners who have continuous contact with children, such as people working in schools and community health services, can have a leading role in recognizing, responding to, and supporting maltreated children. Health professionals such as nurses, midwives, doctors, dentists, and social workers should be urged to 'observe' a child's appearance and behaviour, and look out for any physical or emotional signs of abuse. If a health professional suspects possible maltreatment they should seek an explanation from the child in an open and sensitive way and consult an expert such as a community paediatrician. They should also try to gather information from other agencies and the child should be seen again at some point in the future. However, if there is compelling evidence to suggest that a child is being abused, they should be immediately referred to social services.
2. NICE (2009) has produced guidance that is intended to encourage healthcare professionals to think holistically when a child presents, so that they think about what they see, hear and any other information they receive to help them build up a picture. For example, if maltreatment is suspected, they may need to look at the whole child, gather relevant information from other sources, discuss the case with a senior colleague, and review the child. The guidance is intended to ensure that children who need help get it early in order to prevent further harm, and to enable additional support services to be provided to families where needed.
3. All individual professionals need to recognize that maltreatment is often part of children's lives in households that are also affected by poverty, substance abuse, mental health problems, physical disability, stress, or other forms of violence, which can add significantly to the adverse effects of the child's maltreatment. Enhancing the prospects for healthy development in the lives of maltreated children therefore requires attention to enhancing opportunities for positive, non-violent family and peer interactions.

4. As there are strong associations between child maltreatment and parental mental health conditions or substance misuse, there is a need for professionals to consider the welfare of children when dealing with these problems in adults. These problems, more recently described by the term ‘new morbidity’, have always existed but only recently has their full extent been recognized. More inter-agency collaboration between mental health agencies and those dealing with substance misuse is required.
5. Practitioners need to build on what we already know, to get a better grasp on how abused or neglected children are faring. Child health and development surveys already contain multiple indicators of child wellbeing that could be adapted to suit the purposes of child welfare and child protective service agencies. The challenge now is to develop strategies and resources that can select from these indicators and incorporate them into the routine data collection processes that support agencies’ casework, decision-making, and programme development. However, new measures will need to be developed for evaluating positive outcomes and family strengths.
6. Practitioners frequently have different understandings of what constitutes child abuse and neglect and find it difficult to decide at what point a referral should be made (Horwath, 2005; 2007). In line with the public health approach, identifying the health and psycho-social needs unique to children with a history of various forms of abuse has broad implications for practice, treatment access and planning. Because these needs are often varied and interconnected, an effective inter-agency and multi-professional response is crucial, with the main focus for child maltreatment being primary prevention: preventing new cases of child maltreatment where maltreatment has not yet occurred. Because child maltreatment is a complex behaviour influenced by many factors, it may be easier to intervene to prevent abuse or neglect from developing than to intervene to change behaviours that are already well-established.
7. Given new evidence that trauma in childhood alters the physiology of the brain, it is time for all individual health and social care practitioners to be educated about the full health impact of violence and abuse, and to be trained to explore these issues

either as the true aetiology of their patients' ill-health, or as an underlying potentiating factor that has contributed to it.

8. Nurses, midwives and health visitors as well as specially trained safeguarding nurse practitioners need to develop a trusting relationship with the mother and other family members to promote sensitive, empathic care of their children. They also need to assist mothers to review their own childrearing histories and help them decide how they want to parent their children. School nurses also have a key role in the identification of children who may have been abused or are at risk of abuse.
9. All health organisations should have safeguarding children procedures in place, and a designated or named safeguarding children's nurse whose contact details are known throughout the organisation. Safeguarding children training should be mandatory for all nurses and health workers who may come into contact with children and young people, including ancillary and office staff. This training should be provided on induction, with refreshers at least once a year throughout their employment. These professionals should consider child maltreatment if a child or young person displays a marked change in behaviour or emotional state that is a departure from what would be expected for their age and developmental stage.
10. The availability of appropriate treatments to meet the needs of these children, however, still remains a challenge (Brandon and Thoburn, 2008). Developing effective interventions and services is vital in order to support parents in meeting their children's health and wellbeing needs. Primary prevention efforts could thus be marketed universally, to further reduce the stigma associated with 'parent training': every parent can benefit from parent skills training, not just 'bad' ones.

Policy implications

1. The importance of preventing child maltreatment and thereby its short-term and long-term health and mental health consequences cannot be underestimated. Intervening at an early stage with 'good' parenting programmes may reduce a child's likelihood of developing long-term health problems, and also reduce the public health burden of child maltreatment by preventing future health problems and re-victimization in adulthood with all its negative health consequences.
2. Trust managers should provide access to specialist post-registration safeguarding children education programmes for all professionals working in safeguarding children, as well for as selected professionals who take a lead role in safeguarding children. More and better training is needed to assist professionals in making appropriate use of core assessments and the common assessment framework (CAF) to support abused and neglected children, and to ensure appropriate decisions are made about when to intervene. Within health care, primary-care providers such as family doctors, dentists and A and E (Sidebotham and Biu, 2007) are of particular concern, because they make few referrals to child-protection services despite their ongoing contact with families (Lazenbatt and Freeman, 2006; Flaherty et al, 2008; Woodman et al, 2008).
3. Children's rights as laid out in the UN convention on the rights of the child (UN General Assembly, 1989) provide a framework for understanding child maltreatment as part of a range of violence, harm, and exploitation of children at the individual, institutional, and societal levels. The principles embodied in the UNCRC are concordant with those of medical ethics. The greatest strength of an approach based on the UNCRC is that it provides a legal instrument for implementing policy, accountability, and social justice, all of which enhance public health responses. Incorporation of the principles of the UNCRC into laws, research, public health policy and professional training and practice will result in further progress in the area of child maltreatment.

Research briefing

4. Improving the context of children's and families' lives, for instance in relation to inequalities and housing, good quality childcare, the benefits system and specialist substance misuse, mental health and domestic violence services have the potential to reduce the likelihood of children suffering health and mental health consequences of maltreatment.²
5. A public education campaign is needed to raise awareness of the extent and seriousness of the consequences of any form of child maltreatment, and the importance of reporting it to the appropriate agencies.

² The NSPCC publishes its position statements on a range of topics related to child protection on NSPCC inform: see [NSPCC Policy summaries](#)

References

Agid O., Kohn Y., Lerer B. (2000) Environmental stress and psychiatric illness, *Biomedical Pharmacotherapy* 54: 135–141.

American Professional Society on the Abuse of Children (1998). *Glossary of terms and the interpretations of findings for child sexual abuse evidentiary examinations*. American Professional Society on the Abuse of Children,

Anderson C.M., Teicher M.H., Polcari A., Renshaw P.F. (2002) Abnormal T2 relaxation time in the cerebellar vermis of adults sexually abused in childhood: potential role of the vermis in stress-enhanced risk for drug abuse. *Psycho-neuro-endocrinology*, 27 (1-2):231-44.

Bacchus L., Mezey G., Bewley S. (2002) Women's perceptions and experiences of routine enquiry for domestic violence in a maternity service. *British Journal of Obstetrics and Gynaecology*, 109: 9-16.

Brandon M, Thoburn J. (2008) Safeguarding children in the UK: a longitudinal study of services to children suffering or likely to suffer significant harm. *Child Family Social Work*; 13: 365–77.

Bremner, J.D., Narayan, M., Anderson, E.R., Staib, L.H., Miller, H.L., Charney, D.S. (2000) Hippocampal volume reduction in major depression. *American Journal of Psychiatry*; 157:115–118.

Briere, J. (1996) *Trauma symptoms checklist for children (TSCC): Professional manual*, Psychological Assessment Resources: Odessa, Florida.

Carlson, B. E. (2000). Children exposed to intimate partner violence: Research findings and implications for intervention. *Trauma, Violence, and Abuse*, 1(4):321- 340.

Carr, A. (ed) (2000) *'What Works with Children and Adolescents?' - A Critical Review of Psychological Interventions with Children, Adolescents and their Families*. London: Brunner-Routledge.

Cawson, P., Wattam, C, Brooker, S and Kelly, G (2000) *Child Maltreatment in the United Kingdom: A Study of the Prevalence of Child Abuse and Neglect*. London: National Society for the Prevention of Cruelty to Children.

Cawson P. (2002) *Child Maltreatment in the Family: The experience of a national sample of young people*. London: National Society for the Prevention of Cruelty to Children.

Chester D.L., Jose R.M., Aldiyami E., King H., Moiemmen N.S. (2006) Non-accidental burns in children—are we neglecting neglect? *Burns*; 32: 222–28.

Cicchetti, D., and Rogosch, F.A. (2001) The impact of child maltreatment and psychopathology upon neuroendocrine functioning. *Development and Psychopathology*, 13: 783-804.

Coid, J., Petrukevitch, A., Feder, G. (2001) Relation between childhood sexual and physical abuse and risk of re-victimisation in women: a cross-sectional survey. *Lancet*, 358: 450-454.

Connolly A., Katz V.I., Bash K.L. (1997) Trauma and pregnancy. *American Journal of Perinatology*, 14 (6): 331-336.

Conway, E.E. (1998). Non-accidental head injury in infants: The shaken baby syndrome revisited. *Paediatric Annals*, 27 (10): 677-690.

Crittenden, P. (1998) Child Neglect: Causes and Contributions. In H. Dubowitz (ed) (1999) *Neglected Children: Research, Practice and Policy*. Thousand Oaks: Sage.

Cuffe, S.P., Addy, C.L., Garrison, C.Z., Waller, J.L., Jackson, K.L., McKeown, R.E., Chilappagari, S. (1998). Prevalence of PTSD in a community sample of older adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 37, 147–154. doi:10.1097/00004583-199802000-00006.

Darves-Bornoz, J.M., Lepine, J.P., Choquet, M., Berger, C., Degiovanni, A., and Gaillard, P. (1998). Predictive factors of chronic Post-traumatic stress disorder in rape victims. *European Psychiatry*, 12: 281–287. doi:10.1016/S0924-9338(98)80045-X.

De Bellis MD. (2005) The psychobiology of neglect. *Child Maltreatment*, 10 (2):150-72.

De Bellis, M., and Thomas, L. (2003) Biologic findings of post-traumatic stress disorder and child maltreatment. *Current Psychiatry Reports*, 5: 108-117.

Edwards, V. J., Holden, G.W., Felitti, V. J. (2003) Relationship between multiple forms of childhood maltreatment and adult mental health in community respondents: results from the adverse childhood experiences study. *American Journal of Psychiatry*, 160:1453-1460.

Flaherty E.G., Sege R.D., Hurley T.P. (2008) Translating child abuse research into action *Paediatrics*; 122 (suppl 1): S1–5.

Dubowitz, H., Papas M.S., Black M.M., and Starr, R.H. (2002) Child Neglect: Outcomes in High-Risk Urban Preschoolers, *Paediatrics*, Vol. 109 No. 6, pp. 1100-1107.

Duncan, R.D. (1999) Maltreatment by Parents and Peers: The Relationship between Child Abuse, Bully Victimization, and Psychological Distress, *Child Maltreatment*, Vol. 4, No. 1: 45-55.

Felitti, V. J. (2002) The relationship of adverse childhood experiences to adult health: Turning gold into lead. *Zeitschrift für Psychosomatische Medizin und Psychotherapie* 48(4): 359-369. Available online at: http://www.acestudy.org/files/Gold_into_Lead_-_Germany1-02_c_Graphs.pdf [accessed 28 January 2010].

Felitti, V. J., Anda, R. F., Nordenberg, D., Williamson, D. F., Spitz, A. M., Edwards, V. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*, 14:245-258.

Fergusson, D.M., Horwood, J., and Lynskey, M.T. (1996) Childhood sexual abuse and psychiatric disorder in young adulthood: II. Psychiatric outcomes of childhood sexual abuse. *Journal of the American Academy of Child and Adolescent Psychiatry*, 34:1365–1374.

Flaherty, E.G., Sege R.D, Hurley T.P. (2008) Effect of early childhood adversity on health, *Archives of Paediatrics and Adolescent Medicine*, 160: 1232-1238.

Finkelhor D. (2008) *Childhood victimization. Violence, crime and abuse in the lives of young people*. Oxford: Oxford University Press.

Finkelhor, D. and Baron, L. (1986) *High risk children*. In *A sourcebook of child sexual abuse*. D. Finkelhor, S. Arajii, L. Baron et al, eds. Beverly Hills, CA: Sage, pp. 60–88.

Garnefski, N. and Arends,E. (1998) Sexual abuse and adolescent maladjustment: Differences between male and female victims. *Journal of Adolescence*, 21 1, pp. 99–107.

Garbarino J. (1986) Troubled youth, troubled families: the dynamics of adolescent maltreatment. In D Cichetti and V Carlson (eds) *Child Maltreatment: Theory and Research on the Causes and Consequences of Child Abuse and Neglect*. Cambridge: Cambridge University Press.

Gelles, R.J. (1998). The youngest victims: Violence toward children. In R.K. Bergen (Ed.), *Issues in intimate violence: (pp. 5-24) Thousand Oaks, CA: Sage: National Research Council*.

Gilbert, R., Kemp, A., Thoburn, J., Sidebotham, P., Radford, L., Glaser, D., MacMillan, H.L. (2008) Recognizing and responding to child maltreatment, *Lancet*, DOI:10.1016/S0140-6736(08)61707-9.

Glaser, D. (2001) Child Abuse and Neglect and the Brain – A Review. *Journal of Child Psychology and Psychiatry and Allied Disciplines* 41 (1): 97-116.

Glaser D, Prior V, Lynch M. (2001) *Emotional abuse and emotional neglect: antecedents, operational definitions and consequences*, York: British Association for the Study and Prevention of Child Abuse and Neglect.

Goodall E. and Lumley T. (2007) *'Not seen and not heard' – child abuse: a guide for donors and funders*, New Philanthropies Capital: London.

Hecht, D.B., and Hansen, D.J. (2001). The environment of child maltreatment: Contextual factors and the development of psychopathology. *Aggression and Violent Behavior*, 6: 433-457.

Heger A., Ticson L., Velasquez O., Bernier R. (2002) Children referred for possible sexual abuse: medical findings in 2384 children. *Child Abuse Neglect*, 26: 645–59.

Heim, C. and Nemeroff, C.B. (2001) The role of childhood trauma in the neurobiology of mood and anxiety disorders: preclinical and clinical studies, *Biological Psychiatry* 49:1023–1039.

Hester, M., Pearson, C., Harwin, N., Doble, J., Fisher, U., Hendry, E., Barnardo's, University of Bristol, School for Policy Studies, NSPCC, Department of Health (DoH) (1998) *Making an impact: children and domestic violence*. Ilford, Essex, Barnardo's.

Home Office. (2003) *British Crime Survey*, Home Office: London.

Horwitz, S. (2003) Reliability of the Services Assessment for Children and Adolescents, *Psychiatric Services*, Vol. 52 No. 8.

Horwath, J. (2005) Is This Child Neglect? The Influence of Differences in Perceptions of Child Neglect on Social Work Practice. In J. Taylor and B. Daniel (eds) *Child Neglect: Practice Issues for Health and Social Care* (73-96). London and Philadelphia: Jessica Kingsley Publishers.

Horwath J. (2007) The missing assessment domain: personal, professional and organisational factors influencing professional judgements when identifying and referring child neglect. *British Journal of Social Work*; 37:1285–303.

Hosking, G.D.C. and Walsh, I.R. (2005) *The WAVE Report 2005: Violence and What to Do about It*, WAVE Trust:UK.

Humphreys C. (2000a) *Social Work, Domestic Violence and Child Protection: Challenging Practice*, Policy Press.

Humphreys C. and Mullender A. (2003) Working with Domestic Violence. In Horwath, J. and Shardlow, S. (eds) *Making Links: Assessment and Roles Across Social Work Specialisms*, Lyme Regis: Russell House Publishing.

Humphreys C. and Stanley N. (2005) Multi-agency and Multi-disciplinary work: Barriers and opportunities. In Humphreys, C. and Stanley, N. (eds) (2005) *Child Protection and Domestic Violence: Directions for Good Practice*, London, Jessica Kingsley Publications.

Humphreys C. (2005) Relevant evidence for practice working with domestic violence and child abuse. In Humphreys, C. and Stanley, N. (eds) (2005) *Child Protection and Domestic Violence: Directions for Good Practice*, London, Jessica Kingsley Publications

Humphreys C., Thiara R.K., Skamballis A. and Mullender A. (2006) *Talking about Domestic Abuse: A Photo Activity Workbook to Develop Communication Between Mothers and Young People*, London, Jessica Kingsley.

Jasinski J.L. (2004) Pregnancy and domestic violence: a review of the literature. *Trauma Violence Abuse*, 5(1): 47-64.

Research briefing

Johnson, R., Rew, L., and Sternglanz, R. W. (2006) The relationship between childhood sexual abuse and sexual health practices of homeless adolescents. *Adolescence*, 41(162):221-234.

Jones, D. and Ramchandani, P. (1999) *Child Sexual Abuse - Informing Practice from Research*. Oxford: Radcliffe Medical Press.

Kelley, B. T., Thornberry, T. P., & Smith, C. A. (1997) *In the wake of childhood maltreatment*. Washington, DC: National Institute of Justice. Available at: www.ncjrs.gov/pdffiles1/165257.pdf [Accessed 22 February 2010]

Kendall-Tackett, K.A. (2000) *Treating the lifetime health effects of childhood victimization*. Kingston, NJ: Civic Research Institute Inc.

Kendall-Tackett K.A. (2001) Physiological correlates of childhood abuse: chronic hyper-arousal in PTSD, depression, and irritable bowel syndrome. *Child Abuse Neglect*, 24(6):799-810.

Kendall-Tackett K.A. (2003) *Treating the lifetime health effects of childhood victimization* Kingston, NJ: Civic Research Institute Inc.

Kendler, Kenneth S. and Gardner, Charles O. (1998) Twin Studies of Adult Psychiatric and Substance Dependence Disorders: Are They Biased By Differences in the Environmental Experiences of Monozygotic and Dizygotic Twins in Childhood and Adolescence, *Psychological Medicine*, 28: 625-33.

Kolko, D.J. (2002) Child physical abuse. In: Myers J.E.B., Berliner L., Briere J., Hendrix C.T., Reid T.A., Jenny C.A., editors. *The APSAC Handbook on Child Maltreatment*. Thousand Oaks, CA: Sage Publications, Inc.

Lanktree C.B., Gilbert A.M., Briere J., et al. (2008) Multi-informant assessment of maltreated children: convergent and discriminant validity of the TSCC and TSCYC. *Child Abuse Neglect*, 32: 621-25.

Lazenbatt A. and Freeman R. (2006) Recognizing and reporting child physical abuse: a cross-sectional survey of primary health care professionals, *Journal of Advanced Nursing*, 56, 3: 227- 237.

Laming, Lord (2003) *The Victoria Climbié Inquiry*. London: HMSO.

McCauley, J., Kern, D.E., Kolodner, K., Dill, L., Schroeder, A.F., DeChant, H.K., Ryden, J., Derogatis, and L.R., Bass, E.B. (1997) *Clinical characteristics of women with a history of childhood abuse: unhealed wounds*. *JAMA* May 7;277(17):1362–1368

McLeer, S.V., Deblinger, E.B., Henry, D., and Orvaschel, H. (1992). Sexually abused children at high risk for post-traumatic stress disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*, 31, 875–879. doi:10.1097/00004583- 199209000-00015.

Research briefing

McGee C. (2000) Childhood experiences of domestic violence, *Adoption and Fostering*, 20: 8-15.

McVeigh C., Hughes K., Bellis M., Reed E., Ashton J., Syed Q. (2005) *Violent Britain. People, prevention and public health*. Liverpool: John Moores University.

Manly J.T., Cicchetti D., Barnett D. (1994) The impact of subtype, frequency, chronicity, and severity of child maltreatment on social competence and behaviour problems. *Developmental Psychopathology*, 6:121–143.

Mezey G.C. and Bewley S. (1997) Domestic violence and pregnancy. *British Medical Journal*, 314: 1295.

Mullen, P. E., Martin, J. L., Anderson, J. C., Romans, S. E., and Herbison, P. (1993). Child sexual abuse and mental health in adult life. *British Journal of Psychiatry*, 163: 721–732.

Mullender A., Hague G., Imam U., Kelly L., Malos E. and Regan, L. (2003) *Children's Perspectives on Domestic Violence*, London, Sage.

Mullender, A. (2005) What children tell us: "He said he was going to kill our mum", in Humphreys, C. and Stanley, N. (eds) *Child Protection and Domestic Violence: Directions for Good Practice* (2005) London, Jessica Kingsley.

NICE (2009) *When to suspect abuse – clinical guideline*, National Collaborating Centre for Women's and Children's Health: RCOG Press: London. Accessible online at: www.nice.org.uk/nicemedia/pdf/CG89FullGuideline.pdf [Accessed 22 February 2010]

Noll J.G., Horowitz L.A., Bonanno G.A., Trickett P.K., and Putnam F.W. (2003) Revictimization and self-harm in females who experienced childhood sexual abuse: results from a prospective study, *Journal of Interpersonal Violence*, Dec 18(12):1452-71.

Noll, J.G., Trickett, P.K., Susman, E. and Putman, F.W. (2006) Sleep disturbances and child sexual abuse, *Journal of Paediatric Psychology*, 31(5):469-480.

Oates, M. (2003) Suicide: the leading cause of maternal death, *The British Journal of Psychiatry*, 183: 279-281

Royal College of Paediatrics and Child Health (2007) *Paediatric Forensic Examination in Relation to Possible Child Sexual Abuse*, London: RCPCH, Faculty of Forensic and Legal Medicine.

Radford, L. and Hester, M. (2006) *Mothering Through Domestic Violence*. London: Jessica Kingsley.

Read, J. (1998) Child abuse and the severity of disturbance among adult psychiatric inpatients, *Child Abuse & Neglect*, Vol. 22, No. 5:359–368.

Rosen, L. N., and Martin, L. (1996) Impact of childhood abuse history on psychological symptoms among male and female soldiers in the US army. *Child Abuse & Neglect*, 20: 1149–1160.

Schore, A. N. (2003) Early relational trauma, disorganized attachment, and the development of a predisposition to violence. In M. F. Solomon and D. J. Siegel (Eds.), *Healing trauma: Attachment, mind, body, and brain*. New York, NY: Norton.

Sidebotham P. (2007) Fatal Child Maltreatment. In: Sidebotham P, Fleming P, eds. *Unexpected death in childhood*. Chichester: Wiley: 75–94.

Sidebotham P. and Golding J. (2001) Child maltreatment in the 'children of the nineties'- a longitudinal study of parental risk factors, *Child Abuse and Neglect*, 25, 9: 1177-1200.

Sidebotham P, Biu, T., Goldsworthy, L. (2007) Child protection procedures in emergency departments. *Emergency Medicine Journal*; 24: 831–35.

Silvern, L., Karyl, J., Waelde, L., Hodges, W. F., Starek, J., Heidt, E., and Min, K. (1995). Retrospective reports of parental partner abuse: Relationships to depression, trauma symptoms and self-esteem among college students. *Journal of Family Violence*, 10: 177-202.

Silverman, A.B., Reinherz, H.Z., and Giaconia, R.M. (1996) The long-term sequelae of child and adolescent abuse: A longitudinal community study. *Child Abuse and Neglect*, 20(8): 709-723.

Singer, M.I., Miller, D.B., Guo, S., Slovak, K., and Frierson, T. (1998). *The mental health consequences of children's exposure to violence*. Cleveland, OH: Cayahoga County Community Mental Health Research Institute, Mandel School of Applied Social Sciences, Case Western Reserve University.

Springer, K.W., Sheridan, J., Kuo, D., and Carnes, M. (2007) Long-term physical and mental health consequences of childhood physical abuse: Results from a large population-based sample of men and women. *Child Abuse and Neglect*, 31: 517-530.

Teicher, M.D. (2000) Wounds that time won't heal: The neurobiology of child abuse. *Cerebrum: The Dana Forum on brain science*, 2(4): 50-67.

Teicher M.H., Dumont N.L., Ito Y., Vaituzis C., Giedd J.N., Andersen S.L. (2004) Childhood neglect is associated with reduced corpus callosum area. *Biological Psychiatry*; 56(2):80-5.

Trickett, P.K., and Putnam, F.W. (1998) Developmental consequences of child sexual abuse. In P.K. Trickett and C. J. Schellenbach (Eds.), *Violence against children in the family and the community*:39-56. Washington, DC: American Psychological Association.

Vinchon M., Defoort-Dhellemmes S., Desurmont M., Dhellemmes P. (2005) Accidental and non-accidental head injuries in infants: a prospective study. *Journal of Neurosurgery*; 102: 380–84.

Vullimay A.P., Sullivan R. (2000) Reporting child abuse: paediatricians experiences with the child protection system. *Child Abuse Neglect*; 24: 1461–70.

Walby S. and Allen J. (2004) *Domestic Violence, Sexual Assault and Stalking: Findings from the British Crime Survey*, Home Office Research Study 276, London: Home Office.

Research briefing

Wallace, H. (1996) *Family violence: Legal, medical, and social perspectives*. Needham Heights, MA: Allyn and Bacon.

Walrath, C.M., Ybarra, M.L., Sheehan, A.K., Holden, E.W., Burns, B.J. (2003) Impact of maltreatment on children served in community mental health programs, *Journal of Emotional and Behavioral Disorders*: 14: 73-81.

Weniger G., Lange C., Sachsse U., Irle E. (2008) Amygdala and hippocampal volumes and cognition in adult survivors of childhood abuse with dissociative disorders. *Acta Psychiatrica Scandinavica*, 118:281–90.

Widom, C. S. (1999). Post-traumatic stress disorder in abused and neglected children grown up. *The American Journal of Psychiatry*, 156:1223–1229.

Widom, C.S. and Maxfield, M.G. (2001) *An update on the 'cycle of violence'*. Washington, DC: National Institute of Justice.

Woodman J., Pitt M., Wentz R., Taylor B., Hodes D., and Gilbert R.E. (2008) *Performance of screening tests for child physical abuse in accident and emergency departments*, Health Technology Assessment Programme Monograph Series.

NSPCC

Weston House
42 Curtain Road
London EC2A 3NH

Tel: 020 7825 2500

Fax: 020 7825 2525

www.nspcc.org.uk

Registered charity numbers 216401 and SC037717.

